

CLAIMS

We claim:

Sub 1 1. A method of regulating osteoclast activity, the method comprising causing a soluble RANK to bind RANKL.

2. The method of claim 1, wherein the soluble RANK is encoded by a DNA selected from the group consisting of:

(a) a DNA encoding a protein having an amino acid sequence as set forth in SEQ ID NO:2, wherein the protein has an amino terminus selected from the group consisting of an amino acid between amino acid 1 and amino acid 33, inclusive, of SEQ ID NO:2, and a carboxy terminus selected from the group consisting an amino acid between amino acid 196 and amino acid 616, inclusive;

(b) a DNA encoding a protein having an amino acid sequence as set forth in SEQ ID NO:3, wherein the protein has an amino terminus selected from the group consisting of an amino acid between amino acid 1 and amino acid 30, inclusive, of SEQ ID NO:3, and a carboxy terminus selected from the group consisting an amino acid between amino acid 197 and amino acid 625, inclusive;

(c) DNA molecules capable of hybridization to the DNA of (a) or (b) under stringent conditions, and which encode RANK polypeptides that bind RANKL; and

(d) DNA molecules encoding fragments of proteins encoded by the DNA of (a), (b) or (c), wherein the fragments of RANK polypeptides bind RANKL.

3. The method of claim 2, wherein the RANK is at least about 80% identical in amino acid sequence to native RANK

4. The method of claim 3, wherein the RANK further comprises a polypeptide selected from the group consisting of an immunoglobulin Fc domain, an immunoglobulin Fc mutein, a FLAGTM tag, a peptide comprising at least about 6 His residues, a leucine zipper, and combinations thereof.

Sub 5 5. A method of ameliorating effects of excess bone loss, comprising administering a soluble RANK polypeptide composition to an individual at risk for excess bone loss, and allowing the soluble RANK to bind RANKL and inhibit binding thereof to cells expressing RANK.

Sub 6
6. The method of claim 5, wherein the individual is at risk from or suffers from a condition selected from the group consisting of osteoporosis, Pagett's disease, and bone cancer, and cancers associated with hypercalcemia.

a
7. The method of claim 5, wherein the soluble RANK is encoded by a DNA selected from the group consisting of:

(a) a DNA encoding a protein having an amino acid sequence as set forth in SEQ ID NO:2, wherein the protein has an amino terminus selected from the group consisting of an amino acid between amino acid 1 and amino acid 33, inclusive, of SEQ ID NO:2, and a carboxy terminus selected from the group consisting an amino acid between amino acid 196 and amino acid 616, inclusive;

(b) a DNA encoding a protein having an amino acid sequence as set forth in SEQ ID NO:6, wherein the protein has an amino terminus selected from the group consisting of an amino acid between amino acid 1 and amino acid 30, inclusive, of SEQ ID NO:6, and a carboxy terminus selected from the group consisting an amino acid between amino acid 197 and amino acid 625, inclusive;

(c) DNA molecules capable of hybridization to the DNA of (a) or (b) under stringent conditions, and which encode RANK polypeptides that bind RANKL; and

(d) DNA molecules encoding fragments of proteins encoded by the DNA of (a), (b) or (c), wherein the fragments of RANK polypeptides bind RANKL.

8. The method of claim 7, wherein the RANK is at least about 80% identical in amino acid sequence to native RANK

Sub 3
9. The method of claim 8, wherein the RANK further comprises a polypeptide selected from the group consisting of an immunoglobulin Fc domain, an immunoglobulin Fc mutein, a FLAGTM tag, a peptide comprising at least about 6 His residues, a leucine zipper, and combinations thereof.

10. The method of claim 6, wherein the soluble RANK is encoded by a DNA selected from the group consisting of:

(a) a DNA encoding a protein having an amino acid sequence as set forth in SEQ ID NO:2, wherein the protein has an amino terminus selected from the group consisting of an amino acid between amino acid 1 and amino acid 33, inclusive, of SEQ ID NO:2, and a carboxy terminus selected from the group consisting an amino acid between amino acid 196 and amino acid 616, inclusive;

(d) DNA molecules encoding fragments of proteins encoded by the DNA of (a), (b) or (c), wherein the fragments of RANK polypeptides bind RANKL.

12. The method of claim 11, wherein the RANK further comprises a polypeptide selected from the group consisting of an immunoglobulin Fc domain, an immunoglobulin Fc mutein, a FLAGTM tag, a peptide comprising at least about 6 His residues, a leucine zipper, and combinations thereof.